



Integrity ★ Service ★ Excellence

Atomic and Molecular Physics Program

7 March 2012

**Tatjana Curcic
Program Manager
AFOSR/RSE
Air Force Research Laboratory**

Report Documentation Page			Form Approved OMB No. 0704-0188	
<p>Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p>				
1. REPORT DATE 07 MAR 2012	2. REPORT TYPE	3. DATES COVERED 00-00-2012 to 00-00-2012		
4. TITLE AND SUBTITLE Atomic And Molecular Physics Program			5a. CONTRACT NUMBER	
			5b. GRANT NUMBER	
			5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)			5d. PROJECT NUMBER	
			5e. TASK NUMBER	
			5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory,Wright Patterson AFB,OH,45433			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)	
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited				
13. SUPPLEMENTARY NOTES Presented at the Air Force Office of Scientific Research (AFOSR) Spring Review Arlington, VA 5 through 9 March, 2012				
14. ABSTRACT				
15. SUBJECT TERMS				
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 40
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	19a. NAME OF RESPONSIBLE PERSON	



2012 AFOSR SPRING REVIEW



NAME: Tatjana Curcic

BRIEF DESCRIPTION OF PORTFOLIO:

Understanding interactions between atoms, molecules, ions, and radiation.

LIST SUB-AREAS IN PORTFOLIO:

- **Cold Quantum Gases**
 - Strongly-interacting quantum gases
 - Ultracold molecules
 - New phases of matter
 - Non-equilibrium quantum dynamics
- **Quantum Information Science (QIS)**
 - Quantum simulation
 - Quantum communication
 - Quantum metrology, sensing, and imaging
 - Cavity optomechanics



Scientific and Transformational Opportunities



Scientific Opportunities	Transformational Opportunities
Dipolar Matter, Ultracold Molecules	<ul style="list-style-type: none">• Novel phases of matter• Ultracold chemistry
Quantum Memories and Interfaces	<ul style="list-style-type: none">• Long-distance quantum communication
Quantum Simulation	<ul style="list-style-type: none">• High-T_c superconductivity• Novel phases of matter
Quantum Metrology and Sensing	<ul style="list-style-type: none">• Ultra-high-precision clocks• High-resolution, high-sensitivity magnetometry• High-sensitivity gravimetry• Precision inertial navigation in GPS-denied environments
Non-equilibrium Quantum Dynamics	<ul style="list-style-type: none">• Dynamic control of materials• Efficient optical devices



Outline



- **Quantum Simulation, Strongly-Interacting Quantum Gases**
 - **Bosons: Markus Greiner, Harvard (MURI)**
 - **Algorithmic Cooling in Quantum Gases**
Waseem Bakr, *et al*, *Nature* **480**, 500 (2011)
 - **Quantum Magnetism**
Jonathan Simon, *et al*, *Nature* **472**, 307 (2011)
 - **Fermions: Martin Zwierlein, MIT (PECASE, MURI)**
 - **Evolution of Fermi Pairing from 3D to 2D**
Ariel T. Sommer, *et al*, *Phys. Rev. Lett.* **108**, 045302 (2012)
 - **Spin Transport in a Strongly-Interacting Fermi Gas**
Ariel Sommer, *et al*, *Nature* **472**, 201 (2011)
 - **Thermodynamics of a Unitary Fermi Gas: Superfluid Lambda Transition**
Mark J.H. Ku, *et al*, *Science* (in print); K. Van Houcke, *et al* (submitted to *Nature Physics*)
- **Dipolar Matter: Benjamin Lev, UIUC/Stanford (YIP)**
 - **Dy BEC, and First Dipolar Degenerate Fermi Gas**
Mingwu Lu, *et al*, *Phys. Rev. Lett.* **107**, 190401 (2011)
- **Quantum Metrology: Till Rosenband, NIST**
 - **Coherent Drive Spectroscopy**
D.B. Hume, *et al*, *Phys. Rev. Lett.* **107**, 243902 (2011)




Strongly-correlated quantum gases



Optical lattices

Atoms interact strongly on lattice sites
Superfluid - Mott insulator transition





Synthetic matter, quantum simulations


Electrons in a crystal lattice




Atoms in an optical lattice




Strongly correlated materials



High Tc superconductors



Quantum magnets



Quasi-low
dimensional
materials







Quantum magnetism




Ising model:
*quantum phase transition
from paramagnet to
antiferromagnet*



**Quantum gas
microscope**



**Algorithmic
cooling**




**Quantum
magnetism**



Quantum Gas Microscope




Waseem S. Bakr, et al, *Nature* 463, 74 (2009)




Many new possibilities for

- Detection
- Manipulation
- Cooling
- Creating new systems






Interaction induced orbital excitation blockade



Axial vibrational
excitation

$$|n_{\text{ground}}, n_{\text{excited}}\rangle$$






Orbital excitation blockade operations



$$n_{tot} = n+m$$





Algorithmic cooling



Random occupation




Filter operations
 $4 \rightarrow 3$, $3 \rightarrow 2$, $2 \rightarrow 1$



Mott insulator




Superfluid





Algorithmic cooling






Quantum Magnetism



Quantum simulation of an antiferromagnetic Ising spin chain




$h_x = 0$: classical first order phase transition

Finite h_x : quantum phase transition, second order



Tilted Hubbard Model and Mapping to Spin Model



realizes constraint drives transition

$$H = J \sum_i [S_z^i S_z^{i+1} - \underbrace{(1 - \tilde{D})}_{h_z} S_z^i - \underbrace{2^{3/2} \tilde{t}}_{h_x} S_x^i]$$



$$\tilde{D} = E - U$$

E: energy difference per lattice site, or lattice tilt
U: onsite interaction



Adiabatic transition to the AF state

Jonathan Simon, et al, *Nature* 472, 307 (2011)



- High magnetic field:
spins align with field

- Low magnetic field:
antiferromagnetic
interactions
dominate, producing
Neel order

Modulation spectroscopy:
turn double occupancy
into single occupancy

$$\text{[Yellow Square]} = \uparrow \quad \text{[Black Square]} = \downarrow$$




DISTRIBUTION A: Approved for public release; distribution is unlimited.

Direct spin imaging
preliminary






Selecting homogeneous domain



- 6 adjacent sites that transition simultaneously (RMS shift 6 Hz)
- Compare to 10%-90% width (105 Hz) and transverse field (28 Hz)




Transitioning at highest and lowest energy many-body state



Jonathan Simon, et al, *Nature* **472**, 307 (2011)



Direct measurement of Neel order parameter






Outline


- **Quantum Simulation, Strongly-Interacting Quantum Gases**
 - Bosons: Markus Greiner, Harvard (MURI)
 - Algorithmic Cooling in Quantum Gases
Waseem Bakr, *et al*, *Nature* **480**, 500 (2011)
 - Quantum Magnetism
Jonathan Simon, *et al*, *Nature* **472**, 307 (2011)
 - Fermions: Martin Zwierlein, MIT (PECASE, MURI)
 - Evolution of Fermi Pairing from 3D to 2D
Ariel T. Sommer, *et al*, *Phys. Rev. Lett.* **108**, 045302 (2012)
 - Spin Transport in a Strongly-Interacting Fermi Gas
Ariel Sommer, *et al*, *Nature* **472**, 201 (2011)
 - Thermodynamics of a Unitary Fermi Gas: Superfluid Lambda Transition
Mark J.H. Ku, *et al*, *Science* (in print); K. Van Houcke, *et al* (submitted to *Nature Physics*)
- Dipolar Matter: Benjamin Lev, UIUC/Stanford (YIP)
 - Dy BEC, and First Dipolar Degenerate Fermi Gas
Mingwu Lu, *et al*, *Phys. Rev. Lett.* **107**, 190401 (2011)
- Quantum Metrology: Till Rosenband, NIST
 - Coherent Drive Spectroscopy
D.B. Hume, *et al*, *Phys. Rev. Lett.* **107**, 243902 (2011)



Strongly Interacting Fermi Gases in coupled quasi-2D layers



**High- T_c Superconductor
with stacks of CuO planes**



**Stacks of 2D coupled
fermionic superfluids**

- Access physics of layered superconductors
- Evolution of Fermion Pairing from 3D to 2D
- Berezinskii-Kosterlitz-Thouless transition in deep 2D limit




Evolution of Fermion Pairing from 3D to 2D



Ariel T. Sommer, et al, *Phys. Rev. Lett.* **108**, 045302 (2012)


Density of States: 3D $\sim \sqrt{\epsilon}$
2D *const.*

Direct consequence:
Always a bound state in 2D




$$V_0 = 2 E_R$$
$$V_0 = 5 E_R$$
$$V_0 = 6 E_R$$
$$V_0 = 10 E_R$$
$$V_0 = 12 E_R$$
$$V_0 = 19 E_R$$
$$V_0 = 20 E_R$$

RF spectroscopy:






Evolution of Fermion Pairing from 3D to 2D



- Observed fermion pairing from 3D to 2D
 - Are those pairs superfluid?
 - Study coherence, thermodynamics, rotation...
 - What is T_c as a function of interlayer tunneling?

Towards a recipe for high T_c










Spin Transport – “Little Fermi Collider” (LFC)



Ariel Sommer, et al, *Nature* 472, 201 (2011)



Spin Diffusion vs. Temperature






Thermodynamics of the Strongly-Interacting Fermi Gas




Revealing the Superfluid Lambda Transition in a Fermi Gas

Mark J.H. Ku, *et al*, *Science* (in print)



“Feynman diagrams versus Feynman quantum emulator”

K. Van Houcke, *et al* (submitted to *Nature Physics*)





Outline



- Quantum Simulation, Strongly-Interacting Quantum Gases
 - Bosons: Markus Greiner, Harvard (MURI)
 - Algorithmic Cooling in Quantum Gases
Waseem Bakr, *et al*, *Nature* **480**, 500 (2011)
 - Quantum Magnetism
Jonathan Simon, *et al*, *Nature* **472**, 307 (2011)
 - Fermions: Martin Zwierlein, MIT (PECASE, MURI)
 - Evolution of Fermi Pairing from 3D to 2D
Ariel T. Sommer, *et al*, *Phys. Rev. Lett.* **108**, 045302 (2012)
 - Spin Transport in a Strongly-Interacting Fermi Gas
Ariel Sommer, *et al*, *Nature* **472**, 201 (2011)
 - Thermodynamics of a Unitary Fermi Gas: Superfluid Lambda Transition
Mark J.H. Ku, *et al*, *Science* (in print); K. Van Houcke, *et al* (submitted to *Nature Physics*)
- Dipolar Matter: Benjamin Lev, UIUC/Stanford (YIP)
 - Dy BEC, and First Dipolar Degenerate Fermi Gas
Mingwu Lu, *et al*, *Phys. Rev. Lett.* **107**, 190401 (2011)
- Quantum Metrology: Till Rosenband, NIST
 - Coherent Drive Spectroscopy
D.B. Hume, *et al*, *Phys. Rev. Lett.* **107**, 243902 (2011)




Ultracold Dipolar Physics



- Exploit strong, long-range r^{-3} , and anisotropic dipole-dipole interaction

$$V_d(r) = \frac{d^2}{r^3} (1 - 3\cos^2 \theta)$$





Ultracold dipolar Bose and Fermi gases: Exotic quantum phases




Traps

Bosons

Trap distortions, instabilities
spinors




Quantum ferrofluids



^{52}Cr
Pfau, et al. '08

Fermions



Quantum liquid crystals



nematic

Lattices


Density waves and supersolids



E. Fradkin, S. A. Kivelson, &
V. Oganesyan, Science '07






Laser cooling Dysprosium



$\mu = 10 \mu_B$
 Rb: $200x \mu^2 m$
 Cr: $9x \mu^2 m$

Er MOT: McClelland & Hanssen PRL '06



Five isotopes, 3 bosons ($I=0$), 2 fermions ($I=5/2$)





Strongly Dipolar Dy BEC



M. Lu, N. Burdick, S.-H. Youn, and B. Lev, *Phys. Rev. Lett.* 107 190401 (2011)

BECs of ^{164}Dy and ^{162}Dy



DISTRIBUTION A: Approved for public release; distribution is unlimited.






Evaporative cooling of all isotopes


Including identical fermions! $T/T_F = 1.3$ in crossed ODT




Single beam ODT



Temperature



Number



Ultracold dipolar Bose-Fermi mixtures!

Universal dipolar scattering?


Bohn, Cavagnero, Ticknor, New J. Phys. '09



First dipolar degenerate Fermi gas!




^{161}Dy Sympathetically cooled with ^{161}Dy to $T/T_F < 0.3$




Oblate initial trap

Fermionic ^{161}Dy time-of-flight expansion

Single shot: 6×10^3 atoms



Average of three shots






Outline



- **Quantum Simulation, Strongly-Interacting Quantum Gases**
 - **Bosons: Markus Greiner, Harvard (MURI)**
 - **Algorithmic Cooling in Quantum Gases**
Waseem Bakr, *et al*, *Nature* **480**, 500 (2011)
 - **Quantum Magnetism**
Jonathan Simon, *et al*, *Nature* **472**, 307 (2011)
 - **Fermions: Martin Zwierlein, MIT (PECASE, MURI)**
 - **Evolution of Fermi Pairing from 3D to 2D**
Ariel T. Sommer, *et al*, *Phys. Rev. Lett.* **108**, 045302 (2012)
 - **Spin Transport in a Strongly-Interacting Fermi Gas**
Ariel Sommer, *et al*, *Nature* **472**, 201 (2011)
 - **Thermodynamics of a Unitary Fermi Gas: Superfluid Lambda Transition**
Mark J.H. Ku, *et al*, *Science* (in print); K. Van Houcke, *et al* (submitted to *Nature Physics*)
- **Dipolar Matter: Benjamin Lev, UIUC/Stanford (YIP)**
 - **Dy BEC, and First Dipolar Degenerate Fermi Gas**
Mingwu Lu, *et al*, *Phys. Rev. Lett.* **107**, 190401 (2011)
- **Quantum Metrology: Till Rosenband, NIST**
 - **Coherent Drive Spectroscopy**
D.B. Hume, *et al*, *Phys. Rev. Lett.* **107**, 243902 (2011)




Motivation: Al⁺ clock accuracy





Quantum Logic Spectroscopy








Al⁺ quantum-logic spectroscopy



1. Cool to motional ground-state with Mg⁺ (Raman cooling)
2. Depending on Al⁺ clock state, add one vibrational quantum via 1S_0 - 3P_1
3. Detect vibrational quantum with Mg⁺

QND



P.O. Schmidt, et al.
Science **309**, 749 (2005)

D. B. Hume, et al.
PRL **99**, 120502 (2007)




Al^+ coherent-drive spectroscopy





Mg⁺ Doppler

1. Cool to motional ground state with Mg⁺ (Raman cooling)
drive coherent motion
2. Depending on Al⁺ clock state, add one vibrational quantum via ${}^1\text{S}_0 - {}^3\text{P}_1$
3. Detect ~~vibrational quantum~~ with Mg⁺ coherent motion



QND



D. B. Hume, et al.
PRL 107, 243902 (2011)

${}^3\text{P}_1$ (300 μs) Measure
 ${}^3\text{P}_0$ (21 s) quantum state
w/o scattering
66.6% photons ${}^{27}\text{Al}^+$
 ${}^1\text{S}_0$ Simplified lasers
(no ground-state cooling)
Slower






Al⁺ coherent-drive spectroscopy



D. B. Hume, et al., PRL107, 243902 (2011)


a





Quantum Jumps

Quantum jumps between clock states (1S_0 and 3P_0)



93% state-detection fidelity within 80 ms

with quantum logic:
99% within 10 ms

- Coherent drive detection rate could be improved by higher modulation amplitude or photon collection efficiency
- Can be generalized for more than one Al^+ ion



Interactions with Other Agencies



Agency/Group	POC	Scientific Area
ARO	Peter Reynolds	Cold Quantum Gases (CQG)
	Paul Baker	
	TR Govindan	Quantum Information Science (QIS)
	Rich Hammond	Ultrafast/Ultraintense Phenomena (UUP)
ONR	Charles Clark	CQG, QIS
	Ralph Wachter	QIS
DARPA	Jamil Abo-Shaeer	CQG, QIS
	Jag Shah	QIS
	Matt Goodman	QIS
NSF	Wendell Hill	CQG, QIS, UUP
DoE	Jeff Krause	CQG, UUP
IARPA	Michael Mandelberg	QIS
QISCOG	>20 program managers from ~10 agencies/institutions	QIS

Thank you